



**COUNCIL OF
THE EUROPEAN UNION**

Brussels, 29 November 2012

17032/12

**AGRILEG 181
DENLEG 116
VETER 80**

COVER NOTE

from:	European Commission
date of receipt:	26 November 2012
to:	General Secretariat of the Council
No Cion doc.:	D018801/11
Subject:	Commission Regulation amending Regulation (EC) No 2073/2005 as regards microbiological criteria for sprouts and the sampling rules for poultry carcasses and fresh poultry meat

Delegations will find attached Commission document D018801/11.

Encl.: D018801/11



Brussels, **XXX**
SANCO/10038/2012 Rev. 2
(POOL/G4/2012/10038/10038R2-
EN.doc) D018801/11
[...](2012) **XXX** draft

COMMISSION REGULATION (EU) No .../..

of **XXX**

**amending Regulation (EC) No 2073/2005 as regards microbiological criteria for sprouts
and the sampling rules for poultry carcasses and fresh poultry meat**

(Text with EEA relevance)

EN

EN

COMMISSION REGULATION (EU) No .../..**of XXX****amending Regulation (EC) No 2073/2005 as regards microbiological criteria for sprouts
and the sampling rules for poultry carcasses and fresh poultry meat**

(Text with EEA relevance)

THE EUROPEAN COMMISSION,

Having regard to the Treaty on the functioning of the European Union,

Having regard to Regulation (EC) No 852/2004 of the European Parliament and of the Council of 29 April 2004 on the hygiene of foodstuffs¹, and in particular Article 4(4) thereof,

Whereas:

- (1) Regulation (EC) No 852/2004 lays down general rules for food business operators on the hygiene of foodstuffs, taking particular account of procedures based on the application of hazard analysis and critical control point (HACCP) principles. Article 4 of that Regulation provides that food business operators are to adopt specific hygiene measures concerning inter alia compliance with microbiological criteria for foodstuffs and sampling and analysis requirements.
- (2) Commission Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs² lays down microbiological criteria for certain micro-organisms and the implementing rules to be complied with by food business operators when implementing the general and specific hygiene measures referred to in Article 4 of Regulation (EC) No 852/2004.
- (3) Chapter 1 of Annex I to Regulation (EC) No 2073/2005 sets out the food safety criteria to be complied with by certain food categories, including sampling plans, analytical reference methods and limits for micro-organisms or their toxins and metabolites. That Chapter lists the food safety criteria for sprouted seeds, as regards Salmonella.
- (4) Following the outbreak of Shiga toxin producing *E.coli* (STEC) in May 2011 in the Union, consumption of sprouts was identified as the most likely origin of the outbreaks.

¹ OJ L 139, 30.4.2004, p. 1

² OJ L 338, 22.12.2005, p. 1.

- (5) On 20 October 2011 the European Food Safety Authority ("EFSA") adopted a Scientific Opinion on the risk posed by Shiga toxin producing *Escherichia coli* and other pathogenic bacteria in seeds and sprouted seeds³. In its Opinion, EFSA concludes that the contamination of dry seeds with bacterial pathogens is the most likely initial source of the sprout-associated outbreaks. In addition, the Opinion states that, due to the high humidity and the favourable temperature during sprouting, bacterial pathogens present on dry seeds can multiply during sprouting and result in a public health risk.
- (6) In its Opinion, EFSA recommends *inter alia* that microbiological criteria should be strengthened as one of the components of a food safety management system for the sprouted seeds production chain. That recommendation concerns the existing microbiological criteria on *Salmonella* for sprouted seeds and the consideration of microbiological criteria on other pathogens. EFSA also states that available data indicates a higher risk for sprouts in comparison with other sprouted seeds.
- (7) In its Opinion, EFSA considers different options for microbiological criteria for pathogenic *E. coli* for seeds: before the start of the production process, during sprouting and in the final product. In that context, EFSA states that the detection and mitigation of a contamination problem earlier in the sprouted seed production chain may have advantages as it avoids contamination being amplified during the full sprouting process. It also acknowledges that testing seeds alone does not allow for the detection of contamination which may come at a later stage in the production process. EFSA therefore concludes that microbiological criteria could be useful during the sprouting process and/or for the final product. When considering a microbiological criterion for the final sprouted seeds, EFSA notes that the time required for the detection methods for pathogenic bacteria combined with the short shelf-life may not allow withdrawing the product in the event of non-compliance. In its opinion EFSA considers that it is currently not possible to evaluate the extent of public health protection provided by specific microbiological criteria for seeds and sprouted seeds. This highlights the need for data collection to conduct quantitative risk assessment. Therefore this criterion should be reviewed taking into account progress in science, technology and methodology, emerging pathogenic microorganisms in foodstuffs and information from risk assessment.
- (8) In order to ensure the protection of public health in the Union and in view of that EFSA Opinion, Commission Regulation (EU) No .../ 2013 on certification requirements for imports into the Union of sprouts and seeds intended for the production of sprouts⁴ and Commission implementing Regulation No .../2013 on traceability requirements for sprouts and seeds intended for the production of sprouts⁵ were adopted.
- (9) In addition to the measures laid down in those acts and considering the potential major health risk posed by the possible presence of pathogens in sprouts, provisions on additional microbiological criteria should be laid down following the EFSA recommendations, in particular on STEC serogroups which are considered of most important public health concern.

³ EFSA Journal 2011;9(11):2424.

⁴ OJ L xx, xx.xx.xxxx, p. xx. (SANCO/10064/2012)

⁵ OJ L xx, xx.xx.xxxx, p..xx. (SANCO/10030/2012)

- (10) Microbiological criteria are one of several control options in food safety and should be used by food business operators as a means to verify implementation of an effective food safety management system. However, due to the low prevalence and the heterogeneous distribution of some pathogens in seeds and sprouted seeds, the statistical limitations of sampling plans, and the lack of information about the application of good agricultural practices in the production of seeds, it is necessary to test all batches of seeds for the presence of the pathogens in cases where food business operators have not put in place food safety management systems including steps that reduce the microbiological risk. If food management systems are in place and if their effectiveness is confirmed by historical data, a reduction of sampling frequency may be considered. Such frequency should, however, never be less than once per month.
- (11) When laying down microbiological criteria for sprouts, flexibility should be provided with regard to the stages of sampling and the type of samples which are to be taken, in order to take into account the diversity of production systems, while maintaining equivalent food safety standards. In particular, it is appropriate to provide for alternatives to the sampling of sprouts in cases where the sampling is technically difficult. The testing of spent irrigation water for pathogenic bacteria has been proposed as an alternative strategy, as it seems to be a good indicator of the types of micro organisms in the sprouts themselves. Due to the uncertainties regarding the sensitivity of this strategy, it is needed that food businesses operators using this alternative establish a sampling plan, including sampling procedures and sampling points of the spent irrigation water.
- (12) Certain STEC serogroups (namely O157, O26, O103, O111, O145 and O104:H4) are recognized to be those causing the most of the Haemolytic Uremic Syndrome (HUS) cases occurring in the EU. Furthermore serotype O104:H4 caused the outbreak in May 2011 in the Union. Therefore microbiological criteria should be considered for these six serogroups. It cannot be excluded that other STEC serogroups may be pathogenic to humans as well. In fact, such STEC may cause less severe forms of disease such as diarrhoea and or bloody diarrhoea or may also cause HUS and therefore represent a hazard for the consumer's health.
- (13) Sprouts should be considered to be ready-to-eat food, as they can be consumed without the need for cooking or other processing, which would otherwise be effective in eliminating or reducing to an acceptable level pathogenic micro-organisms. Food business operators producing sprouts should therefore comply with the food safety criteria for ready-to-eat food laid down in Union legislation, including the sampling of processing areas and equipment as part of their sampling scheme.
- (14) Regulation (EC) No 2160/2003 of the European Parliament and of the Council of 17 November 2003 on the control of *Salmonella* and other specified food-borne zoonotic agents⁶ aims at ensuring that proper and effective measures are taken to detect and control *Salmonella* and other zoonotic agents at all relevant stages of production, processing and distribution in order to reduce their prevalence and the risk they pose to public health.

⁶ OJ L 325, 12.12.2003, p. 1.

- (15) Regulation (EC) No 2160/2003, as amended by Commission Regulation (EU) No 1086/2011⁷, lays down detailed rules on a *Salmonella* food safety criterion for fresh poultry meat. As a consequence of the amendments made to Regulation (EC) No 2160/2003, Regulation (EC) No 2073/2005 was also amended by Regulation (EU) No 1086/2011. However, by that amendment, certain terminological ambiguities were introduced in the text of Regulation (EC) No 2073/2005. Those ambiguities should, in the interest of clarity and consistency of Union legislation, be clarified.
- (16) Regulation (EC) No 2073/2005 should therefore be amended accordingly.
- (17) The measures provided for in this Regulation are in accordance with the opinion of the of the Standing Committee on the Food Chain and Animal Health and neither the European Parliament nor the Council has opposed them,

HAS ADOPTED THIS REGULATION:

Formatted: Level 1

Article 1

Regulation (EC) No 2073/2005 is amended as follows:

- (1) in Article 2, the following point (m) is added:

Formatted: Level 1

"(m) the definition of 'sprouts' in Article 2(a) of Implementing Regulation (EU) No .../2013⁵ on traceability requirements for sprouts and seeds intended for the production of sprouts;

- (2) Annex I is amended in accordance with the Annex to this Regulation.

Formatted: Level 1

Article 2

This Regulation shall enter into force on the twentieth day following that of its publication in the *Official Journal of the European Union*.

It shall apply from 1 July 2013.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

Done at Brussels,

Formatted: Level 1

For the Commission
The President
José Manuel BARROSO

Formatted: Level 1

⁷ OJ L 281, 28.10.2011, p. 7.

ANNEX

Formatted: Level 1

Annex I to Regulation (EC) No 2073/2005 is amended as follows:

(1) Chapter 1 is amended as follows:

- (a) footnote 12 is deleted.
- (b) in row 1.18 the reference to footnote 12 is replaced by the reference to footnote 23.
- (c) the following row 1.29 and the corresponding footnote 22 and 23 are added:

"1.29 Sprouts ⁽²³⁾	Shiga toxin producing <i>E. coli</i> (STEC) O157, O26, O111, O103, O145 and O104:H4	5	0	Absence in 25 grams	CEN/ ISO TS 13136 ⁽²²⁾	Products placed on the market during their shelf-life
----------------------------------	--	---	---	------------------------	---	---

(22) Taking into account the most recent adaptation by the European Union reference laboratory for *Escherichia coli*, including Verotoxigenic *E. coli* (VTEC), for the detection of STEC O104:H4;

(23) Excluding sprouts that have received a treatment effective to eliminate *Salmonella* spp and STEC."

(2) Chapter 3 is amended as follows:

(a) in Section 3.2, the part concerning "Sampling rules for poultry carcasses and fresh poultry meat" is amended as follows:

Formatted: Level 1

(i) the first paragraph is replaced by the following:

"Slaughterhouses shall sample whole poultry carcasses with neck skin for *Salmonella* analyses. Cutting and processing establishments other than those adjacent to a slaughterhouse cutting and processing meat received only from this slaughterhouse, shall also take samples for *Salmonella* analysis. When doing so, they shall give priority to whole poultry carcasses with neck skin, if available, but ensuring that also poultry portions with skin and/or poultry portions without skin or with only a small amount of skin are covered, and that choice shall be risk-based."

(ii) the fourth paragraph is replaced by the following:

Formatted: Level 1

"For the *Salmonella* analyses for fresh poultry meat other than poultry carcasses, five samples of at least 25 g of the same batch shall be collected. The sample taken from poultry portions with skin shall contain skin and a thin surface muscle slice in case the amount of skin is not sufficient to form a sample unit. The sample taken from poultry portions

without skin or with only a small amount of skin shall contain a thin surface muscle slice or slices added to any skin present to make a sufficient sample unit. The slices of meat shall be taken in a way that includes as much as possible of the surface of the meat."

- (b) the following Section 3.3 is added:

Formatted: Level 1

"3.3 Sampling rules for sprouts

For the purposes of this Section, the definition of batch in Article 2(b) of Implementing Regulation (EU) No .../2013⁵ will apply.

A. General rules for sampling and testing

Formatted: Level 1

1. Preliminary testing of the batch of seeds

Food business operators producing sprouts shall carry out a preliminary testing of a representative sample of all batches of seeds. A representative sample shall include at least 0.5% of the weight of the batch of seeds in sub samples of 50g or be selected based on a structured statistically equivalent sampling strategy verified by the competent authority.

For the purposes of performing the preliminary testing, the food business operator must sprout the seeds in the representative sample under the same conditions as the rest of the batch of seeds to be sprouted.

2. Sampling and testing of the sprouts and the spent irrigation water

Formatted: Level 1

Food business operators producing sprouts shall take samples for microbiological testing at the stage where the probability of finding Shiga toxin producing *E. coli* (STEC) and *Salmonella* spp is the highest, in any case not before 48 hours after the start of the sprouting process.

Samples of sprouts shall be analysed according to the requirements in rows 1.18 and 1.29 of Chapter 1.

However, if a food business operator producing sprouts has a sampling plan, including sampling procedures and sampling points of the spent irrigation water, they may replace the sampling requirement under the sampling plans set out in rows 1.18 and 1.29 of Chapter 1 with the analysis of 5 samples of 200 ml of the water that was used for the irrigation of the sprouts.

In that case requirements set out in rows 1.18 and 1.29 of Chapter 1 shall apply to the analysis of the water that was used for the irrigation of the sprouts, with the limit of absence in 200 ml.

When testing a batch of seeds for the first time, food business operators may only place sprouts on the market if the results of the microbiological analysis comply with rows 1.18 and 1.29 of Chapter 1, or the limit of absence in 200 ml if they analyse spent irrigation water.

Formatted: Level 1

3. Sampling frequency

Food business operators producing sprouts shall take samples for microbiological analysis at least once a month at the stage where the probability of finding Shiga toxin producing *E. coli* (STEC) and *Salmonella* spp is the highest, in any case not before 48 hours after the start of the sprouting process.

B. Derogation from the preliminary testing of all batches of seeds set out in point A.1 of this Section

When justified on the basis of the following conditions and authorised by the competent authority, food business operators producing sprouts may be exempted from the sampling set out in point A.1 of this Section:

- (a) the competent authority is satisfied that the food business operator implements a food safety management system in that establishment, which may include steps in the production process, which reduces the microbiological risk; and,
- (b) historical data confirms that during at least 6 consecutive months prior to granting the authorisation, all batches of the different types of sprouts produced in the establishment comply with the food safety criteria set out in rows 1.18 and 1.29 of Chapter 1.